arXiv:1112.0125v1 [math.AG] 1 Dec 2011

CLASSIFIYNG METRICALLY CONICAL RATIONAL SINGULARITIES

HELGE MØLLER PEDERSEN

Abstract. In this paper we determine a complete list of rational surface singularities which have metrically conical bilipschitz type of its inner metric. We achieve this by using the thick-thin decomposition of Birbrair, Neumann and Pichon.

1. Introduction

If (X, p) is an isolated complex surface singularity, then a choice of embedding of (X, p) into $(\mathbb{C}^N, 0)$ determines a metric on X called the inner metric. We define the distance between two points to be the infimum of the length of arcs in X between them, where we measure the arclength in the standard hermitian structure on $\mathbb{C}^N$. It turns out that the bilipschitz equivalence class of the inner metric, does not depend of the embedding, and hence is an intrinsic property of the singularity.

It is well known that the topology of a germ complex singularity is determined by its link, i.e. the homeomorphism type of $M = X \cap S^{N-1}_\varepsilon$ for small enough ε, where S^{N-1}_ε is the sphere centered at the origin of radius ε. Namely a neighborhood of $p \in X$ is homeomorphic to the cone over the link. This is usually referred to as the singularity is topological conical see for example [Mil68]. An interesting question is now, when is a neighborhood of $p \in X$ equipped with the inner metric bilipschitz equivalent to a standard metric cone? In [BF08] Birbrair and Fernandes gave the first examples, were this is not the case, and in [BFN08, BFN09, BFN10] Birbrair, Fernandes and Neumann show that in the general case (X, p) is not metrically conical. They also give an example in [BFN10] of singularities with the same topology having different bilipschitz geometries. Hence the bilipschitz type of (X, p) is not in general determined by the topology of (X, p).

In [BNP11] Birbrair, Neumann and Pichon defines what they call a thick-thin decomposition of (X, p), which is invariant under bilipschitz semi-algebraic homeomorphisms. (X, p) is then metrically conical if and only if it consists of one thick piece and no thin pieces. A consequence of this is, that the link of a metrically conical (X, p) has to be Seifert fibered. The way they get the thick-thin decomposition of (X, p) is by studying a general linear function on (X, p) under a good resolution of (X, p) which resolves a generic pencil of hyperplane sections. The thick pieces then correspond to the intersections of the strict transform of the general liner function, and they assume that the resolution is minimal with the extra condition that no thick pieces are adjacent.

In this article we are going to use the thin-thick decomposition to determine which rational singularities are metrically conical. Remember that a singularity is rational if its geometric genus is 0. This is an analytic property, but Artin showed in [Art66] that wether a singularity is rational or not is completely determined by the topology of $(X, 0)$.

2000 *Mathematics Subject Classification.* 14B05, 32S05, 32S25; 57M99.

Key words and phrases. bilipschitz geometry, normal surface singularity, rational singularity, thick-thin decomposition.

Even more the strict transform of a general linear function is also determined by the topology, since the maximal ideal cycle is equal to the minimal cycle. Hence we can determine which rational singularities that are metrically conical from the topology.

We will define a family of weighted starshaped graphs $G_{n,k,l}$ depending on the integers n, k, l with $n \geq 1$, $k \geq 0$ and $l \geq 1$ and if $k = 0$ then $n > l$. Each $G_{n,k,l}$ is the plumbing diagram for a rational homology sphere Seifert fibered manifold, and we will prove the following result:

Theorem 1.1. *(X, p) is a rational singularity with metrically conical bilipschitz geometry if and only if $G_{n,k,l}$ is a dual resolution graph for a resolution of (X, p) for some n, k, l.*

The family includes some well known graph, for example $G_{2,0,1}$ is A_1 and $G_{1,1,1}$ is D_4. It is already known that these two are metrically conical.

In Section 2 we describe the thick-thin decomposition of rational singularities. In Section 3 we will determine some relations on the multiplicities of the maximal ideal cycle of a metrically conical singularity. In Section 4 we will define the family $G_{n,k,l}$, and show that if (X, p) has dual resolution graph $G_{n,k,l}$, then (X, p) is a metrically conical rational singularity. Finally in section 5 we will show that these are the only singularities which are both rational and metrically conical.

2. Thick-thin decomposition of rational singularities

There exist a decomposition of (X, p) into thick and thin parts, the former essential metrically conical, and the later shrinking faster than linear while approaching the origin. Such a decomposition is essentially unique (for detalis see [BNP11]). To describe the thick-thin decomposition. We consider a good resolution $\pi \colon \widetilde{X} \to X$, i.e. the exceptional divisor consists of smooth curves intersecting transversely at most two at a time, with the following properties:

1. It resolves a generic pencil of hyperplane sections, i.e. the strict transforms of generic members are disjoint. A curve of the exceptional divisor meeting the strict transform is called a *$\mathcal{L}$-curve.*
2. All $\mathcal{L}$-curves are disjoint.
3. π is minimal with respect to the above conditions.

Such a resolution can always be achieved from a minimal good resolution by blowing up a finite number of points. In the case of rational singularities we will later see that no blow ups are needed to resolve the generic pencil of hyperplane sections.

Let $\Gamma(\widetilde{X})$ be the dual resolution graph of the resolution. A *bamboo* is a string of vertices of valence 2 ending in a vertex of valence 1. A $\mathcal{L}$-node is a vertex corresponding to a $\mathcal{L}$-curve. Remove all $\mathcal{L}$-nodes and edges attached to a $\mathcal{L}$-node from $\Gamma(\widetilde{X})$, then the remaining components are called the *Tjurina components.* For each vertex v of $\Gamma(\widetilde{X})$ let E_v be the corresponding curve of the exceptional divisor E. Let $N(E_v)$ be a small closed tubular neighborhood of E_v. For any subgraph Γ' of $\Gamma(\widetilde{X})$ let:

$$N(\Gamma') := \bigcup_{v \in \Gamma'} N(E_v) \text{ and } \mathcal{N}(\Gamma') := \overline{N(\Gamma) - \bigcup_{v \notin \Gamma'} N(E_v)}$$

Let B_ε be the ball of radius ε in $\mathbb{C}^N$ centered at the origin.

Assume ε is sufficiently small such that $\pi^{-1}(X \cap B_\varepsilon) \subset N(\Gamma)$. For each Tjurina component Γ_i $i = 1, \dots, r$ not a bamboo define

$$Z_i := \pi(\mathcal{N}(\Gamma_i)) \cap B_\varepsilon.$$

Let $U_1, \dots, U_s$ be the closures of the connected components of $\pi^{-1}(B_\varepsilon) - \bigcup_1^r \mathcal{N}(\Gamma_i)$ and denote

$$Y_j := \pi(U_j).$$

Notice that Y_i is $\pi(N(\Gamma_v))$ where Γ_v is a $\mathcal{L}$-node and attached bamboos.

Theorem 2.1 ([BNP11])**.** *The minimal thick-thin decomposition of (X, p) is as follows:*

$$(X, o) = \Big(\bigcup_{i=1}^{r} (Z_i, 0) \Big) \bigcup \Big(\bigcup_{j=1}^{s} (Y_j, 0) \Big). \tag{1}$$

Where each Z_i are thin *and each Y_j are the* thick.

No thick pieces can be adjacent by assumption (2) on out resolution, so if there are more than one thick piece then there will also be at least on thin piece. Using the thick-thin decomposition one gets the following description of metrically conical singularities which is Corollary 7.11 of [BNP11]. By a node we mean a vertex of valence greater than 2 or an $\mathcal{L}$-node.

Proposition 2.2. *(X, p) is metrically conical if and only if there is no thin pieces. Alternatively, there is only one node in Γ and it is the unique $\mathcal{L}$-node.*

This implies that the link is Seifert fibered, and the graph is starshaped.

If $f \colon (X, 0) \to (\mathbb{C}, 0)$ is an analytic function, we get a principal divisor on $\widetilde{X}$ by considering $f \circ \pi$, and a decomposition $(f \circ \pi) = (f)_\Gamma + \mathrm{St}(f)$ where $(f)_\Gamma$ is the part supported on the exceptional divisor, and $\mathrm{St}(f)$ is the part supported on strict transform. The set $\mathcal{Z}_{an} = \{(f)_\Gamma | f \in m_{X,0}\}$ is ordered and has a unique minimal element Z_{max} called *the maximal ideal cycle*. Since $(f \circ \pi) \cdot E_i = 0$ for all E_i curves of the exceptional divisor, $(f)_\Gamma \cdot E_i \leq 0$ for all i, hence if we define $\mathcal{Z}_{top} = \{Z \neq 0 \text{ effective cycle } | Z \cdot E_i \leq 0 \ \forall i\}$ it is clear that $\mathcal{Z}_{an} \subseteq \mathcal{Z}_{top}$. Now $\mathcal{Z}_{top}$ is also a ordered set with a unique minimal element Z_{min} called *the minimal or fundamental cycle*, and it is clear that Z_{min} is a topological candidate for Z_{max}, for more details see for example [Ném99]. For rational singularities we have the following from [Art66]:

Theorem 2.3. *If X is a rational singularity, then $\mathcal{Z}_{an} = \mathcal{Z}_{top}$, and hence for rational singularities $Z_{max} = Z_{min}$.*

What is important for the thick-thin decomposition is that $Z_{max} = (l_{gen})_\Gamma$ for a generic linear function $l_{gen} \colon (X, 0) \to (\mathbb{C}, 0)$. This means we can determine the the thick-thin decomposition of a rational singularity from Z_{min}, provided the resolution resolves a generic pencil of hyperplane sections. For this we have the following:

Proposition 2.4. *Let $\pi \colon \widetilde{X} \to X$ be a resolution of a rational singularity (X, p). Then π resolves a generic pencil of hyperplane sections of (X, p).*

Proof. We will follow the spirit of Némethi‘s proof of Theorem 2.3 in [Ném99]. By a *cut* associated to a curve E_i of the exceptional divisor, we mean a non compact divisor C_i of $\widetilde{X}$ such that $C_i \cdot E_i = 1$ and $C_i \cdot E_j = 0$ for $j \neq i$.

Given $a_i, b_i \in E_i$ such that $a_i, b_i \notin E_j$ for $j \neq i$ and $a_i \neq b_i$, we choose cuts $C_i^{a_i}, C_i^{b_i}$ such that $C_i^{a_i} \cap E_i = \{a_i\}$ and $C_i^{b_i} \cap E_i = \{b_i\}$ for all i. Let $m_i = Z_{min} \cdot E_i \leq 0$, and define the divisors $D_a = Z_{min} + \sum (-m_i) C_i^{a_i}$ and $D_b = Z_{min} + \sum (-m_i) C_i^{b_i}$. Using the argument from Némethi’s proof of Theorem 2.3, there exist holomorphic functions f_a and f_b such that $(f_a \circ \pi) = D_a$ and $(f_b \circ \pi) = D_b$. Hence $\mathrm{St}(f_a) = \sum (-m_i) C_i^{a_i}$ and $\mathrm{St}(f_b) = \sum (-m_i) C_i^{b_i}$. Since $Z_{min} = Z_{max}$ it follows that f_a and

f_b general linear functions, and $f_a + \lambda f_b$ for $\lambda \in \mathbb{C}P^1$ defines a generic pencil of hyperplane sections, which is resolved by π. □

Corollary 2.5. *If* (X, p) *is a rational singularity, then the thick-thin decomposition is determined by* Z_{min}.

To find Z_{min} we are going to use the following algorithm due to Laufer's:

Algorithm 2.6. *Let* (X, p) *be a normal surface singularity, and let* $E = \sum E_i$ *be the exceptional divisor of a resolution of* (X, p).

1. *First let* $Z_1 = E$.
2. *If there exist a* E_i *such that* $Z_l \cdot E_i > 0$ *then let* $Z_{l+1} = Z_l + E_i$.
3. *If* $Z_l \cdot E_i \leq 0$ *for all* i*, then the algorithm stops and* $Z_{min} = Z_l$

Laufer proved in [Lau72] that the algorithm terminates and that it gives Z_{min}.

3. MULTIPLICITIES OF Z_{max} SUPPORTING A METRICALLY CONICAL SINGULARITY

Let (X, p) be a normal surface singularity. In this section we assume that the minimal resolution graph of (X, p) is starshaped as on Figure 1.



Figure 1. ,

Let $\pi\colon \widetilde{X} \to X$ be the minimal good resolution and $\pi^{-1}(0) = E \cup \bigcup E_{ij}$, where E is the central curve and E_{ij} is the j'th curve on the i'th bamboo. In particular $E^2 = -a$ and $E_{ij}^2 = -e_{ij}$. Denote by $[a_1; a_2; \ldots; a_l]$ the continued fraction

$$a_1 - \cfrac{1}{a_2 - \cfrac{1}{a_3 - \ldots}}. \tag{2}$$

We define the integers p_i, q_i, p_{ij} and q_{ij} where $i = 1, 2, \ldots, n$ and $j = 1, 2, \ldots, k_i$ as $p_i/q_i = [e_{i1}, e_{i2}, \ldots, e_{ik_i}]$ and $p_{ij}/q_{ij} = [e_{ij}, e_{i(j+1)}, \ldots, e_{ik_i}]$, notice $p_i = p_{i1}$ and $q_i = q_{i1}$.

Lemma 3.1. *Let* $Z = mE + \sum_{i=1}^{n} \sum_{j=1}^{k_i} m_{ij} E_{ij} \in \mathcal{Z}_{top}$. *Then* $Z \cdot E_{ij} = 0$ *for all* i, j *if and only if* $m = m_{ik_i} p_i$ *for all* i *and* $m_{ij} = m_{ik_i} p_{i(j+1)}$ *for all* i, j.

Proof. $Z \cdot E_{ij} = 0$ for all i, j ii and only if the multiplicities satisfy the following set of equations:

$$m_{ij} e_{ij} = m_{i(j-1)} + m_{i(j+1)} \quad \text{for } 1 < j < k_i - 1 \tag{3}$$

$$m_{ik_i} e_{ik_i} = m_{i(k_i-1)} \tag{4}$$

$$m_{i1} e_{i1} = m + m_{i2}. \tag{5}$$

We want to see that $m_{ij} = m_{ik_i}p_{i(j+1)}$ if and only if these equations are satisfied. We do this by induction in j starting with $j = k_i - 1$. Using (4), we get

$$m_{i(k_i-1)} = m_{ik_i}e_{ik_i} = m_{ik_i}[e_{ik_i}] = m_{ik_i}p_{ik_i}. \tag{6}$$

Assume that $m_{i(j+1)} = m_{ik_i}p_{i(j+2)}$. Then from (3) we get that

$$m_{ij} = m_{i(j+1)}e_{i(j+1)} - m_{i(j+2)} = m_{ik_i}p_{i(j+2)}e_{i(j+1)} - m_{ik_i}p_{i(j+3)}. \tag{7}$$

But

$$[e_{i(j+1)}; e_{i(j+2)}; \dots; e_{ik_i}] = e_{i(j+1)} - \frac{1}{[e_{i(j+2)}; \dots; e_{ik_i}]} = e_{i(j+1)} - \frac{q_{i(j+2)}}{p_{i(j+2)}}, \tag{8}$$

and $q_{i(j+2)} = p_{i(j+3)}$, hence

$$[e_{i(j+1)}; e_{i(j+2)}; \dots; e_{ik_i}] = \frac{p_{i(j+2)}e_{i(j+1)} - p_{i(j+3)}}{p_{i(j+2)}}. \tag{9}$$

Combining equation (7) and (9) gives that $m_{ij} = m_{ik_i}p_{i(j+1)}$. If we consider m as m_{i0} we also get that $m = m_{ik_i}p_i$. The other direction follows from multiplying (9) by m_{ik_i}. ☐

Proposition 3.2. *If $Z_{max} = Z_{min}$ and (X, p) is metrically conical, then $m = \operatorname{lcm}_i(p_i)$.*

Proof. If (X, p) is metrically conical, then $Z_{max} \cdot E_{ij} = 0$ for all i, j. Since from Lemma 3.1 $m = m_{ik_i}p_i$ for all i, then there exist an l such that $m = l \operatorname{lcm} p_i$. Assume that $l \neq 1$. It follows that $l \mid m_{ik_i}$ for all i since $m_{ik_i} = m/p_i = l \operatorname{lcm} p_i/p_i$, and since $m_{ik_i} \mid m_{ij}$ for all i, j we have that $l \mid m_{ij}$ for all i, j. We can then define a divisor $Z = (m/l)E + \sum_{i=1}^{n}\sum_{j=1}^{k_i}(m_{ij}/l)E_{ij}$. Since $Z = \frac{1}{l}Z_{min}$ it implies $Z \in \mathcal{Z}_{top}$. But $Z < Z_{min}$, hence we get a contradiction. ☐

4. The resolution graphs $G_{n,k,l}$

Let start by defining a family of weighted graphs $G_{n,k,l}$ depending on integers $n \geq 1$, $k \geq 0$ and $l \geq 1$. $G_{n,k,l}$ is starshaped, with its central node having weight $-(n+1)$. It has $n+2$ bamboos as follows:

- n bamboos $\Gamma_1, \dots, \Gamma_n$ of lengths lk where all vertices have weight -2.
- One bamboo Γ_A of length k, where all but the last vertex have weight -2, and the last vertex i.e. the vertex of valence one has weight $-(l+1)$.
- And one bamboo Γ_B of length l where every vertex has weight -2 except the vertex adjacent to the node which has weight $-(k+1)$.

The graph $G_{n,k,l}$ is represented on Figure 2.



Figure 2.

When $k \neq 0$ it is not hard to see that the intersection matrix associated with $G_{n,k,l}$ is negative definite, and that the graph is reduced and hence define a Seifert fibered manifold, which is a singularity link. If $k = 0$ then the graph is not reduced, it is the following graph:



By blowing down -1 vertices we get the reduce graph, which then consist of only one vertex with weight $l - (n+1)$. So $G_{n,0,l}$ has negative definite intersection form if and only if $l \leq n$. Since the case $n = l$ is a smooth point, we will in the sequel, restrict to case $l < n$.

Note that the same graphs could correspond to different triples $(n; k; l)$. For example, have $G_{n,k,1} = G_{n,1,k}$. Notice also that while $G_{n,0,l}$ and $G_{n-l+1,0,1}$ are not equal as graphs their reduced graph are the same, and hence define the same Seifert fibered manifold.

Proposition 4.1. *Let (X,p) be a normal surface singularity with a resolution which have dual resolution graph $G_{n,k,l}$ for some n,k,l, then (X,p) is a metrically conical rational surface singularity*

Proof. In proving the theorem. We are going to use the following criterion for being rational by Laufer:

Theorem 4.2 ([Lau72] 4.2)**.** *Let (X,p) be a normal surface singularity and let $E_1, \dots, E_n$ be the irreducible components of the exceptional divisor for a resolution of (X,p). Then (X,p) is rational if and only if, the genus of all the E_i's is 0 and at any step in Algorithm 2.6 for finding Z_{min} one has that $Z_j \cdot E_i \leq 1$*

When we use Algorithm 2.6 on a starshaped graph E will always denote the central curve, notice that if $Z_1 \cdot E \leq 0$ then $Z_1 = Z_{min}$. If this is not the case always continue the algorithm by adding E first, then making sure that all other curves intersect non positive before we add a new E if necessary. We will be using the following notation when going through algorithm 2.6: Let $m_{\tilde{E}}(Z)$ be the multiplicity of the divisor Z at the curve $\tilde{E}$ of the exceptional divisor. Let E be the central curve of a starshaped resolution graph, let Z'_l denote the divisor given by the algorithm after we just added a E to increase $m_E(Z'_l)$ to l, and let Z_l denote divisor such that $m_E(Z_l) = l$ and $Z_l \cdot \tilde{E} \leq 0$ for $\tilde{E} \neq E$. Notice that $Z'_{l+1} = Z_l + E$.

We start with the proof of the case of $k = 0$. Then as we noted before $G_{n,0,l}$ is not reduced, and the reduced graph is a single vertex with weight $l - (n+1)$. A singularity with such a graph is rational, it follows from 4.2 since $Z_{min} = E$. It is also metrically conical by Proposition 2.2.

Now we assume that $k > 0$. To prove that a singularity with dual resolution graph $G_{n,k,l}$ is rational we will run through Algorithm 2.6 and see that the criterion of theorem 4.2 is satisfied. To do this we need to see how the multiplicities along the bamboo changes during the algorithm, for this we need the following results:

Definition 4.3. We define a bamboo with r vertices to be of the type $A(m,i)$ for integers $m, i \geq 1$ and $r \geq m$. If the curves $E_1, \dots, E_{m-1}$ corresponding to the vertices $v_1, \dots, v_{m-1}$ have weight -2, and $E_m \cdot \sum E_j = -i$. $A(m,i)$ is then either:



where there is m vertices along the bamboo, or



where $c \geq 2$ and there may exist or not exist curves after the rightmost dashed line.

Lemma 4.4. *Let Γ be a bamboo of type $A(m, i)$ and assume that E_1 intersects a curve E. Assume as we run through Algorithm 2.6 $m_E(Z)$ continues to increase. Then there are two cases for the multiplicities of Z_{lm} and two cases for the multiplicities of Z_{lm+1}. The two cases for multiplicities along Γ of Z_{lm} are:*

If $l < i+1$ the multiplicities are



If $l = i+1$ the multiplicities are



The two cases for the multiplicities along Γ of Z_{lm+1} are:

If $l < i+1$ the multiplicities are



If $l = i+1$ the multiplicities are



Proof. Starting with $Z'_2 \cdot E_1 = 1$, we have to add E_1. By induction we also have to add the other E_j's for $j < m$, but not E_m, since $E_m^2 + 2E_{m-1} \cdot E_m + E_m \cdot E_{m+1} = E_m \cdot E_{m-1} + E_m \cdot \sum E_j = 1 - i \leq 0$. Next $Z'_3 \cdot E_1 = 1$, so we continuo to add the E_j's. Except this time we do not have to add E_{m-1}, since $2E_{m-1}^2 + 3E_{m-2} \cdot E_{m-1} + E_m \cdot E_{m-1} = 0$. This process continues. So as long as $m_E < m$ we will increase the multiplicity of E_1. When $m_E = m$ then the multiplicities of Z_m along Γ are as follows:



where the curve with $-d$ is the m'th curve. Notice that $Z'_{m+1} \cdot E_1 = -2m + m + 1 + m - 1 = 0$. So the the multiplicities along Γ will not change.

Next $Z'_{m+2} \cdot E_1 = -2m + m + 2 + m - 1 = 1$ hence we need to add E_1. But now $(Z'_{m+2} + E_1) \cdot E_2 = -2(m-1) + m + 1 + m - 2 = 1$ and we need to add E_2. This continue until the m'th curve. Now $(Z'_{m+2} + \sum_{j=1}^{m-1} E_j) \cdot E_m = E_m \cdot (2E_{m-1} + \sum E_j = 2 + E \cdot \sum E_i = 2 - i$. So if $i = 1$ we have to add E_m. But if $i \neq 1$ we do not.

First assume that $i = 1$. Then the configuration of the multiplicities along Γ of Z'_{2m} are going to be:



Now $Z'_{2m} \cdot E_1 = 1$ hence we need to add E_1. We likewise need to add E_2, but we need not add E_3. So the multiplicities of Z'_{2m+1} are



Still $Z'_{2m+1} \cdot E_1 = 1$. So the multiplicities of Z_{2m+1} is:



Now assume that $i > 1$. Then we will continue to increase m_{E_1} until we get to Z'_{2m}. Then the configuration of multiplicities along Γ will be:



again $Z_{2m} \cdot E_1 = 1$. So we add E_1, and get that the multiplicities for Z'_{2m+1} are:



and now $Z'_{2m+1} \cdot E_1 = 0$. So we do not need to add E_1. Notice that we are in the same situation as before we split, i.e. if we have $A(m,2)$ we must now proceed as in the first case, otherwise we must proceed as in the second case. This clearly continues until we get the the situation described in the statement. ☐

Definition 4.5. We define a bamboo with r vertices to be of type $B(m,c,l)$ for integers $m, l > 0$, $c > 1$ and $l < r-1$, if it is of one of these forms:



or



If we are in the second case then $l > 1$.

Lemma 4.6. *Let Γ be a bamboo of type $B(m,c,l)$ and assume that E_1 intersects a curve E. Assume as we run through Algorithm 2.6, that $m_E(Z)$ continues to increase. Then the multiplicities along Γ of Z_k for different values of k are the following:*

The multiplicities of $Z_{i(m+1)-1}$ for $i+1 < c$ are



The multiplicities of $Z_{i(m+1)}$ for $i+1 < c$ are



The multiplicities of $Z_{i(m+1)-jc-1}$ for $(j-1)c - j + 1 \leq i + 1 < jc - j + 1$ and $j < l$ are



where the multiplicities increase by one going from the right until they reach j.

The multiplicities of $Z_{i(m+1)-jc}$ *for* $(j-1)c-j+1 \leq i+1 < jc-j+1$ *and* $j < l$ *are*



where the multiplicities increase by one going from the right until they reach j.

The multiplicities of $Z_{l(m(c-1)-1)-1}$ *are*



The multiplicities of $Z_{l(m(c-1)-1)}$ *are*



We continues the proof of Proposition 4.1 by noticing that $\Gamma_1, \ldots, \Gamma_n$ are all of type $A(kl,1)$, Γ_A is of type $A(k,l)$ and Γ_B is of type$B(k,1,l)$. Denote the curve corresponding to the vertices along Γ_i by E_{ij} $j = 1, \ldots, kl$ starting with the one adjacent to E. Denote the curve corresponding to the vertices along Γ_A by E_j $j = 1, \ldots, k$ starting with the one adjacent to E. Denote the curve corresponding to the vertices along Γ_B by E'_i $j = 1, \ldots, l$ starting with the one adjacent to E. Now

$$Z_1 \cdot E = E^2 + E \cdot E'_1 + E \cdot E_1 + \sum_{i=1}^{n} E \cdot E_{i1} = -(n+1) + 1 + 1 + \sum_{i=1}^{n} 1 = 1. \tag{10}$$

This means we have to add E. Now $Z'_2 \cdot E'_1 = -(k+1)+2 = -k+1 \leq 0$ so we do not have to add E'_1, but unless $k = 1$ using lemma 4.4 we get that Z_2 has multiplicity 2 everywhere except for the end curves and the E'_i's that have multiplicity 1.

$$Z_2 \cdot E = 2E^2 + E \cdot E'_1 + E \cdot E_1 + \sum_{i=1}^{n} E \cdot 2E_{i1} = -2(n+1) + 1 + \sum_{i=1}^{n-1} 2 = 1. \tag{11}$$

We now have to add E again, and unless $k = 2$ we have to add E_1 and the E_{i1}'s.

This process continuous until we reach Z'_k. By that point the multiplicities along Γ_A starts by being 1 at the end curve and continue to increase by 1 until we get to E_1 which has multiplicity k, according to Lamma 4.4. Notice that so far the multiplicity at E is at most k, and hence $Z_s \cdot E'_1 \leq -(k+1) + k = -1$, hence we have not added anymore E'_1's. Now $Z_k \cdot E = 1$ so we add E to get Z'_{k+1}. Now

$$Z'_{k+1} \cdot E_1 = kE_1^2 + (k+1)E \cdot E_1 + (k-1)E_2 \cdot E_2 \tag{12}$$

$$= -2k + k + 1 + k - 1 = 0. \tag{13}$$

If $l = 1$ then

$$Z'_{k+1} \cdot E_1 = kE_{i1}^2 + (k+1)E \cdot E_{i1} + (k-1)E_{i2} \cdot E_{i2} \tag{14}$$

$$= -2k + k + 1 + k - 1 = 0 \tag{15}$$

$$Z'_{k+1} \cdot E'_1 = {E'_1}^2 + (k+1)E \cdot E' = -(k+1) + k + 1 = 0 \tag{16}$$

$$Z_{k+1} \cdot E = (k+1)E^2 + E \cdot E'_1 + kE \cdot E_1 + \sum_{i=1}^{n} E \cdot kE_{i1} \tag{17}$$

$$= -(k+1)(n+1) + 1 + k + \sum_{i=1}^{n} k = -n. \tag{18}$$

Hence we have found Z_{min}. Notice that at each step, we have $Z_l \cdot E_{ij} \leq 1$, $Z_l \cdot E' \leq 1$ and $Z_l \cdot E \leq 1$. So the singularity is rational by Theorem 4.2.

We saw above (in equation (16)) that $Z_{min} \cdot E'_1 = 0$. It is also not hard to see that the bamboos satisfy the equations of Proposition 3.1. Hence since $Z_{min} = Z_{max}$ and E is the only $\mathcal{L}$-curve. The singularity is then metrically conical by Proposition 2.2.

If $l \neq 1$ then

$$Z'_{k+1} \cdot E_1 = kE_{i1}^2 + (k+1)E \cdot E_{i1} + (k-1)E_{i2} \cdot E_{i2} \tag{19}$$

$$= -2k + k + 1 + k = 1 \tag{20}$$

$$Z'_{k+1} \cdot E'_1 = {E'_1}^2 + (k+1)E \cdot E' + E'_2 \cdot E'_1 = -(k+1) + k + 1 + 1 = 1 \tag{21}$$

$$Z_{k+1} \cdot E = (k+1)E^2 + 2E \cdot E'_1 + kE \cdot E_1 + \sum_{i=1}^{n} E \cdot kE_{i1} \tag{22}$$

$$= -(k+1)(n+1) + 2 + k + \sum_{i=1}^{n} (k+1) = 1. \tag{23}$$

Hence the algorithm continues.

Since the Γ_B is of type $B(k, 2, l-1)$ the algorithm will continue until Z_{lk+1} by lemma 4.6. But now since the Γ_i's are of type $A(lk, 1)$ we get that

$$Z_{lk+1} \cdot E = (lk+1)E^2 + lE \cdot E'_1 + (l(k-1)+1)E \cdot E_1 + \sum_{i=1}^{n} E \cdot lkE_{i1} \tag{24}$$

$$= -(lk+1)(n+1) + l + l(k-1) + 1 + \sum_{i=1}^{n} lk = -n. \tag{25}$$

So the algorithm finishes and $Z_{lk+1} = Z_{min}$. Along the way all the intersections have been 1, and hence by Theorem 4.2 (X, p) is rational. It now not hard to see, that the multiplicities along the bamboos satisfies proposition 3.1. Hence E is the only $\mathcal{L}$-curve and (X, p) is metrically conical. ☐

5. Classification of Metrically Conical Rational Singularities

Theorem 5.1. *Let (X, p) be a rational metrically conical singularity, then there exists n, k and l such that $G_{n,k,l}$ is the dual resolution graph for some resolution of (X, p).*

Proof. Let Γ be the dual resolution graph of the minimal good resolution of (X, p).

If Γ only has one vertex with weight $-p$, then $G_{p+1,0,1}$ is Γ blown up at one point, and hence a resolution graph of (X, p). So (X, p) is rational and metrically conical by Proposition 4.1.

The case of Γ being just a single bamboo i.e. when the link is a lens space, is worked out in example 10.2 of [BNP11], and non of those are metrically conical, unless the bamboo only has one vertex. Hence we can from now on assume that the link is Seifert fibered with at least 3 singular fibers.

The way we are going to proceed is by running through Algorithm 2.6. We will then either get to a place where $Z_j \cdot E_i \geq 2$, complete the algorithm but finding a curve other than the central curve has $Z_{min} \cdot E_j < 0$, or if non of these two cases happens, show that $\Gamma = G_{n,k,l}$ for some n, k and l.

We will use the same notation as in section 3 and assume that Γ is the one in Figure 1. We will also reuse the notation from the proof of Proposition 4.1 about divisors. So recall that by $m_{\widetilde{E}}(Z)$ we mean the multiplicity of the divisor Z at the curve $\widetilde{E}$. That Z_i is the candidate for Z_{min} gotten from the algorithm, with $m_E(Z_i) = i$ and $Z_i \cdot \widetilde{E} = 0$ for any curve $\widetilde{E} \neq E$, remember E is the curve corresponding to the central vertex. That $Z'_i = Z_{i-1} + E$. We will run through the algorithm by starting with Z_1. Then add E if necessary and get Z'_2. We will then not add an other E, before we have reached Z_2. Then continue to add E to get Z'_3 and so on.

First if $a \geq n$ then $Z_{min} = Z_1$, and hence at any end curve E' we will have $Z_{min} \cdot E' < 0$, since the self intersecting number is less than -1. So non of those can be metrically conical.

We can then assume that $a < n$. We start by running though Algorithm 2.6. Now $Z_1 \cdot E = E^2 + \sum_{i=1}^{n} E \cdot E_{i1} = -a + n$, hence if $a < n-1$ then $Z_1 \cdot E > 1$ and by Theorem 4.2 X is not rational.

So we assume that $a = n - 1$. Then $Z_1 \cdot E = 1$ and we add E to get our next candidate Z'_2. Next we look at the intersections of Z'_2 with the E_{i1}'s. There are three different cases:

(1) The i'th bamboo is just a single vertex with self intersection number -2, then $Z'_2 \cdot E_{i1} = 0$.
(2) $E_{i1}^2 < -2$, then $Z'_2 \cdot E_{i1} \leq 0$.
(3) $E_{i1}^2 = -2$ and it intersects a curve different from E then $Z'_2 \cdot E_{i1} = 1$.

Hence only in the third case do we need to add the E_{i1}. So assume that $E_{11}, \ldots, E_{s1}$ has intersection as in the third case, all other E_{ij} have intersection as in the first or second case. Hence Z_2 has multiplicity 2 at E and at $E_{11}, \ldots, E_{s1}$ and maybe further along these bamboos, and multiplicity 1 everywhere else. We get that

$$Z_2 \cdot E = 2E^2 + \sum_{i=1}^{n} Z_2 \cdot E_{i1} = -2(n-1) + 2s + n - s = s + 2 - n. \tag{26}$$

If $s = n$ then $\mathbb{Z}_2 \cdot E = 2$ and (X, p) can not be rational. If $s < n - 1$ then the intersection is less than 0 and $Z_2 = Z_{min}$. But by corollary 3.2 this means that all the p_i's associated to the bamboos has to divide 2 and hence be equal to 2 if (X, p) is metrically conical. If this is the case then $\Gamma = G_{n-2,1,1}$.

So we can assume that Γ is the following:



where the vertices past the dashed line need not exist, and $b > 2$ if there is any vertex with weight $-b_1$. We denote the bamboo starting with a vertex of weight $-b$ by Γ', and denote the curve corresponding the first vertex by E', and the following curves by $E_1, \ldots, E_l$.The other n bamboo we call $\Gamma_1, \ldots, \Gamma_n$, and let $E_{i1}, \ldots, E_{ik_i}$ be the curves corresponding to the vertices along Γ_i starting with E_{i1} adjacent to E. Notice that Γ_i is of type $A(m_i, s_i)$ for some $m_i \geq 1$ and $s_i \geq 1$. Let $m = \min_i\{m_i\}$.

We will achieve the proof by treating four different cases depending on the values of m and b.

First case. Assume that $b > m + 1$ or $b = m + 1$ and E_1 does not exist. We will continue running Algorithm 2.6. Remember that we had gotten to a point where $m_E(Z_2) = m_{E_{i1}}(Z_2) = 2$, also along the i'th bamboo the $m_{E_{ij}}(Z_2) = 2$ for $j < m_i$. The multiplicity of all other curves are one. We also had that $Z_2 \cdot E = 1$, hence we have to add E again. By using Lemma 4.4 we see that we need to tun the algorithm until we reach Z_m. Now assume that $m_1 = m$, we can always reorder the bamboos so this is true. Then the multiplicities along the Γ_1 is $m_{E_{11}}(Z_m) = m$, $m_{E_{12}}(Z_m) = m-1$ and so forth. Now $m_{E_{i1}}(Z_m) = m$ for all i, and we have had no need for adding any E'. $Z_m \cdot E = 1$ so we add E again, we now may have to increase the multiplicities of the E_{i1} for $i > 1$ but $m_{E_{11}}(Z_{m+1}) = m$ and $m_{E'}(Z_{m+1}) = 1$, and hence we get that

$$Z_{m+1} \cdot E \leq -(m+1)n + m + 1 + \sum_{i=2}^{n} m + 1 = 0, \tag{27}$$

and hence $Z_{m+1} = Z_{min}$.

Now if $b > m+2$ then $Z_{min} \cdot E' \leq -b + m + 1 + 1 = -1$ and hence E' is and $\mathcal{L}$-curve, so (X, p) is not metrically conical. If $b = m+2$ and E_1 does not exist the same holds. If $b = m+2$ and E_1 exists then $Z_{min} \cdot E' = 0$ but somewhere along the bamboo starting at E' there is a curve E_i such that $Z_{min} \cdot E_i < 0$. So we can assume that $b = m+1$ and E_1 does not exist.

Now assume that there is a $m_i > m$. Then somewhere along the string of -2 on Γ_i we will have the following situation:



and hence the central of the three curves is a $\mathcal{L}$-curve. So (X, p) is not metrically conical.

Finally if $s_i \neq 1$ for some i, then either E_{im_i} is a $\mathcal{L}$-curve or there is a $\mathcal{L}$-curve further along Γ_i. The later case is when $E_{im_i}^2 = -3$.

Now the only case left in our first case is exactly when $\Gamma = G_{n+1,k,1}$ which are rational metrically conical by Proposition 4.1.

Second case. Assume that $b < m$ or $b = m$ and E_1 exists. As in the first case we continue with Algorithm 2.6. But in this case we come to the situation where we have to increase $m_{E'}(Z_l)$ before we reach Z_{min}. Since if E_1 exist we get to Z'_b before we reach Z_{min}, because $b < m+1$ means we will continue to increase the $m_{E_{i1}}(Z_l)$ according to Lemma 4.4. In that case $Z'_b \cdot E' = -b + b + 1 = 1$ and we have to add E', but now

$$Z'_{b+1} \cdot E = -bn + 2 + \sum_{i=1}^{n} b = 2, \tag{28}$$

and hence (X, p) is not rational. The same happens if E_1 does not exist at Z'_{b+1}.

Third case. Assume that $b = m$ and E_1 do not exist. For X to be metrically conical, $bm_{E'}(Z_{min}) = m_E(Z_{min})$ or else E' is an $\mathcal{L}$-curve. Hence b divides $m_E(Z_{min})$. Assume that $m = m_1$. We again run Algorithm 2.6 as in the other cases, and it will not stop before we reach Z_{2b}. But since $m_i \geq m = b$ we get that $Z_{b+1} \cdot E = -nb + 1 + \sum_{i=1}^{n} b = 1$, so the algorithm continues. Now $Z'_{b+1} \cdot E' = 1$ and so we increase $m_{E'}$ to 2. Since $Z'_{b+1} \cdot E_{11} = -2m + (m+1) + (m-1) = 0$ we do not increase $m_{E_{11}}$. If there is an other $m_i = b$ then $Z_{b+1} \cdot E \leq 0$ and the algorithm stops. Since $Z_{b+1} \cdot E' = -b2 + b + 1 < 0$ we get that E' is a $\mathcal{L}$-curve, and (X, p) is not metrically conical. If not then we have that

$$Z_{b+1} \cdot E = -(b+1)n + 2 + b + \sum_{i=2}^{n} (b+1) = 1, \tag{29}$$

so the algorithm continue.

Now if any of the γ_i for $i > 1$ have $m_i < 2b - 1$, and the algorithm will stop at Z_{m_i+1}. But then $Z_{m_i+1} \cdot E' < 0$ and (X, p) is not metrically conical. So we can assume that $m_i \geq 2b - 1$ for $i > 1$, an thus the algorithm will run at least until Z_{2b}.

We now have to split into two subcases depending on wether Γ_1 is of type $A(m, 1)$ or not.

Subcase of bamboo type $A(m, 1)$. We use the calculation of the multiplicities of Lemma 4.4. If one of the $m_i = 2b - 1$, the algorithm stops. But $Z_{2b} \cdot E_{12} = -2(2b - 2) + 2b - 1 + 2b - 4 = -1$, hence we have that E_{12} is and $\mathcal{L}$-curve and (X, p) is not metrically conical.

So we assume that $m_i \geq 2b$ for all $i > 1$ and continue the algorithm by adding E. Now $Z'_{2b+1} \cdot E' = -b2 + 2b + 1 = 1$ and we add E'. Using Lamma 4.4 we get that, if $m_i > 2b$ for all $i > 1$

$$Z_{2b+1} \cdot E = -n(2b + 1) + 2b + 3 + \sum_{i=2}^{n} (2b + 1) = 2, \tag{30}$$

and (X, p) is not rational according to Theorem 4.2 . If $m_i = 2b$ for two or more i's, the algorithm ends, an (X, p) is not metrically conical, since we that E' is an $\mathcal{L}$-curve.

Hence we can assume that $m_2 = 2b$ and $m_i > 2b$ for $i > 2$ and we get that $Z_{2b+1} \cdot E = 1$. We then add E again. Now $Z'_{2b+2} \cdot E' = -b3 + 2b + 2 < 0$ so we do not need to add E', also $Z'_{2b+2} \cdot E_{11} = -2(2b) + 2b + 2 + 2b - 2 = 0$, so we do not need to add E_{11}, and hence the Algorithm ends. But E' is an $\mathcal{L}$-curve unless $b = 2$, so (X, p) is not metrically conical. If $b = 2$ we will study the multiplicities along Γ_2. If $s_2 = 1$ then the multiplicities along Γ_2 are the following according to Lemma 4.4



and one clearly sees that we have a $\mathcal{L}$ at the end of Γ_2. If $s_2 + 1 = b_{2(2b)} > 2$ then from Proposition 3.1 the multiplicities have to satisfy the equation $2b + 2 = m_{k_2} p$ where $p/q = [2; b_{2(2b)}; b_{2(2b+1)} \ldots; b_{2(k_1)}]$. Now $p/q = \frac{2bp' - (2b-1)q'}{(2b-1) - (2b-2)q'}$, where $p'/q' = [b_{2(2b)}; b_{2(2b+1)} \ldots; b_{2k_2}]$. Hence $2b + 2 = M_{k_2}(2bp' - (2b - 1)q')$. So $p' = \frac{b+1}{m_{k_2} b} + \frac{2b-1}{2b} q'$. Since $q' > 1$ we get that $p' < 2q'$ which contradicts that $b_{2(2b)} \neq 2$.

Subcase of bamboo type not $A(m, 1)$. We use the calculation of the multiplicities of from 4.4, and notice that $m_{E'}(Z_{2b}) = 2$ since $Z'_{2b} \cdot E' = -b2 + 2b = 0$. We get that

$$Z_{2b} \cdot E \leq -2bn + 2 + 2b - 1 + \sum_{i=2}^{n} 2b = 1. \tag{31}$$

If the algorithm ends at this point then (X, p) will not be metrically conical, since we see from 4.4 that E_{11} will be a $\mathcal{L}$-curve. So we can assume that $m_i > 2b - 1$.

So we continue the algorithm. But notice we are now in the same situation as when we split into the subcases depending on the bamboo type of Γ_1, except now it is a bamboo of type $A(m, 2)$ that leads to first subcase. The argument above for the first subcese carries through also in this case, and hence (X, p) will not be metrically conical or not be rational. If we do not end up in the first subcase, then either the algorithm is going to stop before we get to Z_{3b} and we will not have a metrically conical singularity, or we are back in this situation again. Since the bamboo type of Γ_1 is $A(m, j)$ for some j, when $m_E = jm$ we will be forced into the first subcase, and hence (X, p) will never be a metrically conical rational singularity.

Fourth case. Assume that $b = m + 1$ and E_1 exists. In this case as in the third case, the the algorithm will run until Z_m without increasing $m_{E'}$. But since we until this point have had to increase $m_{E_{i1}}$ at each step we still have to increase m_E since $Z_m \cdot E = 1$. After this we get that $Z'_{m+1} \cdot E' = -(m + 1) + m + 1 + 1 = 1$ hence we need to add E'. If any of the other m_i is equal to m, then the algorithm stops, but since now $Z_{m+1} \cdot E' \leq -(m + 1)2 + m + 1 + 2 = -m + 1$ we will have that E' is an $\mathcal{L}$-curve.

So we can assume $m_i > m$ for $i > 1$, and as in the third case we have to separate into different subcases, but this time we have four different cases, depending on the bamboo types of Γ' and Γ_1.

Case $A(m, 1)$ and $B(m, 2, l)$. Assume that Γ_1 is of bamboo type $A(m, 1)$ and Γ' is of bamboo type $B(m, 2, l)$. If the algorithm stops before we reach Z'_{2m} then E' is an $\mathcal{L}$-curve. Using the calculation of multiplicities in Lemma 4.4 and 4.6 we get, since $Z_{2m} \cdot E_{12} = -2(2m - 2) + 2m - 1 + 2m - 4 = -1$, that E_{12} is a $\mathcal{L}$-curve, if $Z_{2m} = Z_{min}$. Hence we assume that $m_i > 2m_1$ for $i > 1$. Now we get that:

$$Z_{2m+1} \cdot E = -n(2m + 1) + 2m + 3 + \sum_{i=2}^{n} m_{E_{i1}}(Z_{2m+1}) \tag{32}$$

$$= 2 - (n - 1)(2m + 1) + \sum_{i=2}^{n} m_{E_{i1}}(Z_{2m+1}). \tag{33}$$

If $m_i > 2m$ for $i > 1$ then $Z_{2m+1} \cdot E = 2$ contradicting that X is rational. If more than one of the $m_i = 2m$ the algorithm stops, and we have that E' is an $\mathcal{L}$-curve, since $Z_{2m+1} \cdot E' \leq -(m + 1)3 + 2m + 1 + 3 = -m + 1$. So assume that $m_2 = 2m$, then we have that $Z_{2m+1} \cdot E = 1$ and we add E once more. But now $Z'_{2m+2} \cdot E_{11} = -2(2m) + 2m + 2 + 2m - 2 = 0$ and $Z'_{2m+2} \cdot E' \leq -(m+1)3 + 2m + 2 + 3 = -m + 2$, hence the algorithm stops. In this case E' is a $\mathcal{L}$-curve unless $m = 2$, but in this case one can look at the multiplicities along Γ_2, and this works out as in the first subcase of the third case. We get that there is a $\mathcal{L}$-curve along Γ_2, hence (X, p) is not metrically conical.

Case $A(m, 1)$ but not $B(m, 2, l)$. Assume that Γ_1 is of bamboo type $A(m, 1)$ and Γ' is not of bamboo type $B(m, 2, l)$. Again if the algorithm stops before Z_{2m} then (X, p) is not metrically conical. If the algorithm stops at Z_{2m}, then E' is a $\mathcal{L}$-curve. Hence we can assume that $m_i > 2m - 1$ for $i > 1$. Next $Z_{2m+1} \cdot E \leq -(2m + 1)2 + 2m + 2 + \sum_{i=2}^{n}(2m + 1) = 1$, if the algorithm stops then we have that E_{11} is a $\mathcal{L}$-curve. So we can assume that $m_i > 2m$ for $i > 1$. Notice that we are back in a situation as before we split according to the bamboo type of Γ_1 and Γ'. But since the bamboo type of Γ_1 is already fixed, we can only move into the cases $A(m, 1)$ and $B(m, 3, l)$ or $A(m, 1)$ and not $B(m, 3, l)$. If we move into the first the argument from that case can then be repeated and (X, p) will not be a metrically conical rational singularity. If we stay in this case then (X, p) is either not metrically conical or we end back here again. But either the algorithm ends which means (X, p) is not metrically conical or we are forced into the case $A(m, 1)$ and $B(m, c, l)$ for some c at some point, which again will mean that (X, p) is not metrically conical or (X, p) is not rational.

Case neither $A(m, 1)$ or $B(m, 2, l)$. Assume that Γ_1 is not of bamboo type $A(m, 1)$ and Γ' is not of bamboo type $B(m, 2, l)$. As in the first case, the algorithm runs until Z'_{2m}. If the algorithm stops there, then E_{11} a $\mathcal{L}$-curve, since $Z_{2m} \cdot E_{11} = -2(2m - 1) + 2m + 2m - 3 = -1$. Hence we can assume that we have to add E once more. But $Z'_{2m+1} \cdot E' = -(m + 1)2 + 2m + 1 + 1 = 0$ and $Z'_{2m+1} \cdot E_{11} = -2(2m - 1) + 2m + 1 + 2m - 3 = 0$ and the algorithm stops. If one of Γ_i's do not have $m_i = 2m$ then there will be a $\mathcal{L}$-curve along it. If $s_1 \neq 2$ then there will be a $\mathcal{L}$-curve along Γ_1. And last if $b_1 \neq 2$ then either E_1 or E_2 is a $\mathcal{L}$-curve, this also implies that E_2 can not exist, since then Γ' would be of type $B(m, 2, l)$. But if non of these things happen, then we are in the case where $\Gamma = G_{n+1,m,2}$.

Case $B(m, 2, l)$ but not $A(m, 1)$. Assume that Γ_1 is not of bamboo type $A(m, 1)$ and Γ' is of bamboo type $B(m, 2, l)$. As the other cases we run the algorithm until we reach Z'_{2m} or (X, p) will not be metrically conical. If the algorithm stops at Z_{2m}, we have that E_{11} is a $\mathcal{L}$-curve. Hence we can assume that, $m_i > 2m - 1$ for $i > 1$. We now notice that we are in the same situation before we divided into the 4 subcases. First if Γ_1 is of bamboo type $A(m, 2)$ or not, second if Γ' is of bamboo type $B(m, 2, 2)$ or not. If we have bamboo type $A(m, 2)$ and not $B(m, 2, 2)$, then we are in a situation as the first subcase, and a similar argument as there shows that (X, p) is not rational or not metrically conical. Likewise if we have bamboo type $A(m, 2)$ and $B(m, 2, 2)$, then we are in a similar situation as the second subcase, and we can reuse that argument. Notice that we will here only run through this case once before going into a situation as in the first subcase. If we do not have bamboo type $A(m, 2)$ but have bamboo type $B(m, 2, 2)$, then we are in a case as the third subcase, but now the positive answer comes if $s_1 = 3$, $m_i = 3m$ for $i > 1$ and E_2 is a -2 curve and an end curve. But this is exactly the case of $\Gamma = G_{n-1,m,3}$. If we have neither bamboo type $A(m, 2)$ nor bamboo type $B(m, 2, 2)$ then we will return to this subcase again. Hence we have to split up in cases and we get that Γ is either $G_{n-1,m,4}$, a singularity which is not metrically conical, or we end up here again. But since the algorithm has to stop, this proves the theorem. Since the only rational singularities we do not find to be non metrically conical, are have $\Gamma = G_{n-1,m,l}$.

□

References

[Art66] Michael Artin. On isolated rational singularities of surfaces. *Amer. J. Math.*, 88:129–136, 1966.

[BF08] Lev Birbrair and Alexandre Fernandes. Inner metric geometry of complex algebraic surfaces with isolated singularities. *Comm. Pure Appl. Math.*, 61(11):1483–1494, 2008.

[BFN08] Lev Birbrair, Alexandre Fernandes, and Walter D. Neumann. Bi-Lipschitz geometry of weighted homogeneous surface singularities. *Math. Ann.*, 342(1):139–144, 2008.

[BFN09] Lev Birbrair, Alexandre Fernandes, and Walter D. Neumann. Bi-Lipschitz geometry of complex surface singularities. *Geom. Dedicata*, 139:259–267, 2009.

[BFN10] Lev Birbrair, Alexandre Fernandes, and Walter D. Neumann. Separating sets, metric tangent cone and applications for complex algebraic germs. *Selecta Math. (N.S.)*, 16(3):377–391, 2010.

[BNP11] Lev Birbrair, Walter D. Neumann, and Anne Pichon. The thick-thin decomposition and the bilipschitz classification of normal surface singularities. *arXiv:1105.3327v3*, 2011.

[Lau72] Henry B. Laufer. On rational singularities. *Amer. J. Math.*, 94:597–608, 1972.

[Mil68] John Milnor. *Singular points of complex hypersurfaces.* Annals of Mathematics Studies, No. 61. Princeton University Press, Princeton, N.J., 1968.

[Ném99] A. Némethi. Five lectures on normal surface singularities. In *Low dimensional topology (Eger, 1996/Budapest, 1998)*, volume 8 of *Bolyai Soc. Math. Stud.*, pages 269–351. János Bolyai Math. Soc., Budapest, 1999. With the assistance of Ágnes Szilárd and Sándor Kovács.

Institut de Mathématiques de Luminy, UMR 6206 CNRS, Case 907, 163 Avenue de Luminy, 13288 Marseille cedex 9, France

E-mail address: `helge@imf.au.dk`